





Leibniz
Universität
Hannover

Systematic analysis of atom interferometric measurements in complex gravitational environments

Michael Werner

Institute of Quantum Optics Hannover

02 April 2025



@ Rencontres de Moriond
2025 — Gravitation


Motivation

Atom interferometers like the **VLBAI in Hannover** are highly accurate sensors for the gravitational field and **span about 10 meters**.

New atom interferometers are planned with **baselines of more than 100 meters**.

And even **atom interferometers in space** have been proposed.

Question: How do you model complex gravitational fields and even general relativistic effects in such quantum systems?



PPN Hamiltonian

Our theory colleagues working on general relativity
Domenico Giulini and Philip Schwartz [3] developed a
Hamiltonian describing:

Two point particles (think of Proton and Electron):

1. Coupled gravitationally to the PPN spacetime
2. Coupled electromagnetically to each other
3. EM field coupling to gravity

(Parametrized) Post-Newtonian (PPN) spacetime

$$g_{\mu\nu} = \begin{pmatrix} -1 - 2\frac{\phi}{c^2} - 2\beta\frac{\phi^2}{c^4} & 0 \\ 0 & \left(1 - 2\gamma\frac{\phi}{c^2}\right)\mathbb{1} \end{pmatrix}$$

PPN Hamiltonian

$$H_{\text{C,final}} = \frac{\mathbf{P}^2}{2M} \left[1 - \frac{1}{Mc^2} \left(\frac{\mathbf{p}_r^2}{2\mu} + \frac{e_1 e_2}{4\pi\epsilon_0 r} \right) \right] + \left[M + \frac{1}{c^2} \left(\frac{\mathbf{p}_r^2}{2\mu} + \frac{e_1 e_2}{4\pi\epsilon_0 r} \right) \right] \phi(\mathbf{R})$$

$$- \frac{\mathbf{P}^4}{8M^3 c^2} + \frac{2\gamma + 1}{2Mc^2} \mathbf{P} \cdot \phi(\mathbf{R}) \mathbf{P} + (2\beta - 1) \frac{M \phi(\mathbf{R})^2}{2c^2} ,$$

$$H_{\text{A,final}} = \frac{{}^{(3)}g_{\mathbf{R}}^{-1}(\mathbf{p}_r, \mathbf{p}_r)}{2\mu} + \frac{e_1 e_2}{4\pi\epsilon_0 \sqrt{{}^{(3)}g_{\mathbf{R}}(\mathbf{r}, \mathbf{r})}}$$

$$- \frac{m_1^3 + m_2^3}{M^3} \frac{\mathbf{p}_r^4}{8\mu^3 c^2} + \frac{e_1 e_2}{4\pi\epsilon_0} \frac{1}{2\mu M c^2} \left(\mathbf{p}_r \cdot \frac{1}{r} \mathbf{p}_r + \mathbf{p}_r \cdot \mathbf{r} \frac{1}{r^3} \mathbf{r} \cdot \mathbf{p}_r \right) ,$$

$$H_{\text{AL,final}} = -\mathbf{d} \cdot \mathbf{E}_{\text{coord.}}^{\perp}(\mathbf{R}) + \frac{1}{2M} \{ \mathbf{P} \cdot [\mathbf{d} \times \mathbf{B}_{\text{coord.}}(\mathbf{R})] + \text{H.c.} \}$$

$$- \frac{m_1 - m_2}{4m_1 m_2} \{ \mathbf{p}_r \cdot [\mathbf{d} \times \mathbf{B}_{\text{coord.}}(\mathbf{R})] + \text{H.c.} \}$$

$$+ \frac{1}{8\mu} (\mathbf{d} \times \mathbf{B}_{\text{coord.}}(\mathbf{R}))^2 + \frac{1}{2\epsilon_0} \int d^3\mathbf{x} \left(1 + (\gamma + 1) \frac{\phi}{c^2} \right) \mathcal{P}_d^{\perp 2}(\mathbf{x}, t) ,$$

$$H_{\text{L,final}} = \frac{\epsilon_0}{2} \int d^3\mathbf{x} \left(1 + (1 - 3\gamma) \frac{\phi}{c^2} \right) \left[\mathbf{E}_{\text{phys.}}^{\perp 2} + c^2 \mathbf{B}_{\text{phys.}}^2 \right]$$

$$= \frac{\epsilon_0}{2} \int d^3\mathbf{x} \sqrt{-g} \left[\mathbf{E}_{\text{phys.}}^{\perp 2} + c^2 \mathbf{B}_{\text{phys.}}^2 \right]$$

What we did with this:

We applied this Hamiltonian to atom interferometers.

Created a Python algorithm that **symbolically** calculates the phase shifts (using “SymPy”).

Made it open source for everyone to use it!

Theoretical model:



Link to the algorithm:




What did we do with this algorithm?

We proposed a novel interferometer geometry, the „CGI“

— **“Co-located Gradiometric Interferometer”** —

as a differential measurement between two interferometers:



(Leading) Phase shift of a CGI

Phase comparison of MZI and SDDI

MZI	SDDI	Phase	Magnitude [rad]	Differential signal
2	2	$NkgT_R^2$	1.4×10^7	0
2	2	$Nkz_0\Gamma_0T_R^2$	20	0
2	2	$Nkv_0\Gamma_0T_R^3$	14	0
$-\frac{7}{6}$	$-\frac{7}{6}$	$Nkg\Gamma_0T_R^4$	14	0
2	0	$\frac{N^2\hbar k^2\Gamma_0T_R^3}{m}$	1.5×10^{-2}	2
-6	-6	$\frac{N\omega_R g^2 T_R^3}{c^2}$	2.3×10^{-9}	0
6	6	$\frac{N\omega_R g v_0 T_R^2}{c^2}$	2.4×10^{-9}	0

$$\Delta\Phi = 2 \frac{N^2 \hbar k_R^2 \Gamma_0 T_R^3}{m}$$

$$= f \cdot \Gamma_0$$

$$f = 2 \frac{N^2 \hbar k_R^2 T_R^3}{m}$$


Gravitational potential: $\phi(z) = \phi_0 + gz - \frac{1}{2}\Gamma_0 z^2 + \mathcal{O}(z^3)$

Ideal gravitational fields

Consider $\phi(z) = \phi_0 + gz - \frac{1}{2}\Gamma_0 z^2$ with a constant gravity gradient.

with a non-trivial $\Gamma(z) = \partial_z^2 \phi(z) =$ Gravitational curvature

Example: Gravitational field of VLBAI Hannover [1, 2]



CGI: Numerical simulation in VLBAI

CGI
Phase

Gravitational
Curvature


$$\Delta h = 0.44 \text{ m}$$

$\Delta\Phi(z_0)$ [mrad]

$$\Delta h = 1.75 \text{ m}$$

z_0 [m]

Intuitively clear: the atoms „average“
gravity along their trajectory.
But how exactly?




Averaging gravity

Where are the atoms „on average“?

Answer: It depends on the averaging process!

$$\|z(t)\|_n = \left(\frac{1}{2T_R} \int_0^{2T_R} |z(t) - z_0|^n dt \right)^{1/n}$$

n	1	2	3	4	5
$\ z(t)\ _n$	$0.66 \Delta h$	$0.73 \Delta h$	$0.77 \Delta h$	$0.79 \Delta h$	$0.82 \Delta h$



Estimator for grav. curvature

We can now define a novel estimator $\hat{\Gamma}(z_0)$ for the grav. curvature:

Step 1: Convert phase to grav. curvature using the scale factor.

$$\hat{\Gamma}(z_0) = \frac{\Delta\Phi(z_0)}{f}$$


Step 2: Shift the height via the **cubic** mean of the trajectory.


$$\hat{\Gamma}(z_0) = \frac{\Delta\Phi(z_0 - \|z(t)\|_3)}{f}$$

Reminder:

$$f = -2 \frac{N^2 \hbar k_R^2 T_R^3}{m}$$


Motivation for the **cubic** mean.





Outlook / Summary

- We introduced the „CGI“ — giving us information about gravitational curvature.
- Gives rise to a novel way to measure gravitational field in large baseline interferometers.
- Possibility to measure the gravitational curvature using more compact devices.





Hammerer group

Thanks for listening!
Any questions?



Gaaloul group

Recent paper:



Link to the algorithm:



Other References:

- **[1]** Schilling et al. „Gravity field modelling for the Hannover 10m atom interferometer“
- **[2]** Lezeik et al. „Understanding the gravitational and magnetic environment of a very long baseline atom interferometer“
- **[3]** Schwartz et al. „Post-Newtonian Hamiltonian description of an atom in a weak gravitational field.“

Finite speed of light (FSL)

There are additional phase shifts, if one includes the FSL into the model!

Example: Two-Photon Bragg transitions would lead to an additional phase

$$\Delta\Phi_{\text{FSL}} = \frac{4\hbar N^2 k^2 T_R}{mc} \left(4gT_R - \nu_0 - \frac{N\hbar k}{m} \right) + \Delta\Phi_0,$$


with a constant shift

$$\Delta\Phi_0 = \frac{2\hbar N^2 k^2}{mc} (2z_L - z_0 - z_U).$$

Lower laser position Upper laser position

Not that problematic,
since constant in time.


FSL with two-photon Bragg transitions



Photon paths of the example (FSL effect exaggerated).

FSL mitigation scheme


The time-dependent FSL phase can be mitigated!




Alter the last AIF pulse by:

$$\Delta_{\text{Det}}(\nu_0, T_R) = 2 \frac{\nu_0 + \frac{\hbar k}{m} - 4gT_R}{\nu_0 + \frac{\hbar k}{m} - gT_R} \frac{\hbar k}{mc}$$

This frequency chirp (~ 100 MHz) nullifies the FSL phase.



Gradiometry in complex fields



$$\hat{\Gamma}_0 = \frac{\Delta g}{\Delta z}$$

Each Mach-Zehnder Interferometer (MZI) measures „its“ g .
Divide change in g over the height difference.

If you want high spatial resolution $\rightarrow$ Make Δz and Δh small
 $\rightarrow$ Higher measurement uncertainty